

## Stop Hypertension With the Acupuncture Research Program (SHARP)

### Results of a Randomized, Controlled Clinical Trial

Eric A. Macklin, Peter M. Wayne, Leslie A. Kalish, Peter Valaskatgis, James Thompson, May C.M. Pian-Smith, Qunhao Zhang, Stephanie Stevens, Christine Goertz, Ronald J. Prineas, Beverly Buczynski, Randall M. Zusman

**Abstract**—Case studies and small trials suggest that acupuncture may effectively treat hypertension, but no large randomized trials have been reported. The Stop Hypertension with the Acupuncture Research Program pilot trial enrolled 192 participants with untreated blood pressure (BP) in the range of 140/90 to 179/109 mm Hg. The design of the trial combined rigorous methodology and adherence to principles of traditional Chinese medicine. Participants were weaned off antihypertensives before enrollment and were then randomly assigned to 3 treatments: individualized traditional Chinese acupuncture, standardized acupuncture at preselected points, or invasive sham acupuncture. Participants received  $\leq 12$  acupuncture treatments over 6 to 8 weeks. During the first 10 weeks after random assignment, BP was monitored every 14 days, and antihypertensives were prescribed if BP exceeded 180/110 mm Hg. The mean BP decrease from baseline to 10 weeks, the primary end point, did not differ significantly between participants randomly assigned to active (individualized and standardized) versus sham acupuncture (systolic BP:  $-3.56$  versus  $-3.84$  mm Hg, respectively; 95% CI for the difference:  $-4.0$  to  $4.6$  mm Hg;  $P=0.90$ ; diastolic BP:  $-4.32$  versus  $-2.81$  mm Hg, 95% CI for the difference:  $-3.6$  to  $0.6$  mm Hg;  $P=0.16$ ). Categorizing participants by age, race, gender, baseline BP, history of antihypertensive use, obesity, or primary traditional Chinese medicine diagnosis did not reveal any subgroups for which the benefits of active acupuncture differed significantly from sham acupuncture. Active acupuncture provided no greater benefit than invasive sham acupuncture in reducing systolic or diastolic BP. (*Hypertension*. 2006;48:838-845.)

**Key Words:** acupuncture ■ blood pressure ■ hypertension ■ randomized clinical trial  
■ traditional Chinese medicine

Hypertension affects  $\approx 700$  million individuals worldwide<sup>1</sup> and is attributable each year for  $>7$  million excess deaths and loss of 64 million disability-adjusted life years.<sup>2</sup> Hypertension in most individuals remains untreated or uncontrolled.<sup>1</sup> Effective control of hypertension is limited by availability, cost, and adverse effects of antihypertensive medications.<sup>3</sup> Modalities of complementary and alternative medicine, including acupuncture, are being used by patients with increasing frequency,<sup>4</sup> but these therapies lack demonstrated efficacy and safety for treating cardiovascular disease and hypertension.<sup>5</sup>

Acupuncture has been used in traditional Chinese medicine (TCM) to treat symptoms related to hypertension for  $>2500$  years.<sup>6</sup> Today, acupuncture is commonly used to treat hypertension in China and the West.<sup>7-9</sup> The efficacy of acupuncture is well supported for treating postoperative dental pain<sup>10</sup> and

nausea<sup>11,12</sup> with few reported adverse effects.<sup>13</sup> Acupuncture has been found effective for treating a number of other acute<sup>14-16</sup> and chronic<sup>17,18</sup> conditions in a growing number of randomized trials, although opinion differs on the role of placebo effects.<sup>19,20</sup> Mechanistic studies have demonstrated effects of acupuncture on the activity and plasma concentrations of blood pressure modulators, including: renin, aldosterone, angiotensin II, norepinephrine, serotonin, enkephalins, and  $\beta$ -endorphins.<sup>21-29</sup> The efficacy of acupuncture for treating hypertension is suggested by a large number of published case series and uncontrolled trials.<sup>22,23,25,30-32</sup> Three randomized trials<sup>33-35</sup> reported significant reductions in BP relative to randomly assigned control groups treated for 4 to 8 weeks, whereas 3 others did not report significant effects of acupuncture relative to control subjects.<sup>36-38</sup> They were all relatively small trials ( $n=10$  to  $68$ ), and all but Yin et al<sup>35</sup> were limited

Received February 22, 2006; first decision March 12, 2006; revision accepted August 8, 2006.

From the New England Research Institutes (E.A.M., L.A.K., S.S.), Watertown, Mass; New England School of Acupuncture (P.M.W., P.V.), Watertown, Mass; Children's Hospital Boston (L.A.K.), Boston, Mass; Massachusetts General Hospital (J.T., M.C.M.P.-S., Q.Z., B.B., R.M.Z.), Harvard Medical School, Boston, Mass; Samueli Institute (C.G.), Alexandria, Va; and the School of Medicine (R.J.P.), Wake Forest University, Winston-Salem, NC.

Correspondence to Eric A. Macklin, New England Research Institutes, Inc, 9 Galen St, Watertown, MA 02472. E-mail EMacklin@NERIScience.com  
© 2006 American Heart Association, Inc.

*Hypertension* is available at <http://www.hypertensionaha.org>

DOI: 10.1161/01.HYP.0000241090.28070.4c

by  $\geq 1$  of the following: incomplete descriptions of methods, incomplete blinding, and lack of peer review.

The Stop Hypertension with the Acupuncture Research Program (SHARP) pilot trial<sup>39</sup> overcomes these limitations by combining rigorous methodology, adherence to principles of TCM acupuncture, and a large sample size. We investigated whether acupuncture administered according to principles of TCM more effectively controlled hypertension than invasive sham acupuncture and whether acupuncture individualized to each participant's TCM diagnosis was more effective than acupuncture at preselected acupuncture points.

## Methods

### Trial Design

The SHARP trial was a prospective, double-blind, randomized, sham-controlled, parallel-group clinical trial testing the efficacy of 6 weeks of twice-weekly TCM acupuncture for treating moderate essential hypertension without use of antihypertensive medications. The complete study design has been published.<sup>39</sup>

The trial protocol was approved by applicable institutional review boards and was monitored by an independent data and safety monitoring board. Prospective study participants were recruited from outpatient clinics, referrals from Boston-area physicians, via advertising in periodicals and on public transportation, and through mass mailings to age-eligible Boston residents. All of the participants provided written informed consent. The eligibility criteria included: (1) essential hypertension with stable blood pressure (BP) after suspension of any antihypertensive medications between 140/90 mm Hg and 179/109 mm Hg over 3 qualifying visits spanning 8 to 31 days; (2) age  $\geq 18$  years; (3) no acupuncture in the previous 6 months; and (4) no medical contraindications to acupuncture or suspension of antihypertensive medications if prescribed previously (including: history of cerebrovascular events, endocrine disorders, and renal insufficiency). Participants were randomly assigned with equal allocation to each of 3 treatments: TCM individualized (IND), TCM standardized (STD), or control (CNTL) acupuncture (Table 1). IND and STD acupuncture, based on TCM principles, were considered active treatment. CNTL acupuncture was an invasive sham control with insertion of acupuncture needles at non-TCM sites. Randomizations were stratified according to use of antihypertensive medications in the previous 6 months and were supplied by a centralized computer just before first treatment.

Participants were weaned off antihypertensives, screened for trial eligibility, and followed for 1 year after random assignment for BP, physical exams, electrocardiograms, quality of life (QoL), and

adverse events at Massachusetts General Hospital. Antihypertensive medications were suspended individually with  $\approx 7$  days between each suspension conditional on BP remaining  $< 180/110$  mm Hg. At each visit, 3 or 5 replicate measurements of seated BP were made after at least a 5-minute rest period using manual mercury sphygmomanometers by trained staff masked to participant random assignment. The BP for a given visit was calculated as the average of the second and third or second through fifth BP measurements depending on whether the second and third measurements of both systolic and diastolic BP were or were not within 6 mm Hg of each other, respectively. Baseline BP was calculated as the average from 3 qualifying BPs measured on separate visits with the first qualifying BP occurring  $\geq 7$  days after suspension of any antihypertensive medications, no 2 qualifying BPs closer than 4 days apart, and no longer than 31 days from first to third qualifying BP. After random assignment, BP was monitored at scheduled visits every 14 days out to the primary end point at week 10 and at 4, 6, 9, and 12 months after random assignment. The same size of arm cuff, determined individually for each participant, was used at all of the visits. At 2 visits during the 6- to 8-week treatment interval, BP was measured immediately before and immediately after treatment. Antihypertensive drugs were initiated between random assignment and week 10 under the following conditions: (1) either systolic or diastolic BP exceeded 180 mm Hg or 110 mm Hg, respectively, at a single visit; (2) either systolic or diastolic BP exceeded 170 mm Hg or 105 mm Hg, respectively, on 2 sequential visits; or (3) an adverse event required treatment to lower BP. Criteria for initiating antihypertensive drugs were relaxed after week 10 with drugs initiated if either systolic or diastolic BP exceeded 140 mm Hg or 90 mm Hg, respectively, or to treat an adverse event.

Participants received acupuncture diagnoses and treatment from a team of licensed acupuncturists in a clinic established at Massachusetts General Hospital. Diagnoses and treatment followed TCM principles as described by Cheng.<sup>40</sup> With full follow-up, treatment consisted of  $\leq 12$  generally twice-weekly 30-minute acupuncture sessions provided over 6 to 8 weeks. At visits 1, 5, 9, and 12, a diagnosing acupuncturist trained in China and with  $\geq 12$  years of clinical experience, masked to random assignments, established each participant's TCM differential diagnosis<sup>6</sup> based on evaluation of history and symptoms and examination of pulses, face, and tongue using a 32-point standardized diagnosis inventory. A separate team of treating acupuncturists, masked to BP, administered the most recently prescribed treatment for IND participants or substituted STD or CNTL acupuncture for participants randomly assigned to those treatment groups. Details of the criteria used for TCM diagnoses, acupuncture point definitions, including depth and duration of needle insertion, needle stimulation, elicitation of de qi (a dull aching and irradiating sensation indicative of effective needle placement), and locations of control points are provided in Kalish et al.<sup>39</sup>

Blood chemistries, lipid profiles, complete blood cell counts, and urinalysis were performed at baseline, 10 weeks, 6 months, and 12 months. QoL data were collected using self-administered Medical Outcomes Study Short Form questionnaires<sup>41</sup> at baseline, 10 weeks, and 12 months. Maintenance of participant masking to their treatment randomization was assessed at 4 months using a self-administered instrument that asked participants to guess their treatment assignment and state their confidence on a 5-point scale.

In addition to changes in BP, adverse events were recorded when participants presented with complaints at any time and based on asking participants if they had been well at each visit. Serious adverse events were recorded as defined by the International Conference on Harmonization Guideline E2A.<sup>42</sup>

### Statistical Analysis

The primary end point was change in systolic BP (SBP) from baseline to week 10. Ten-week change in diastolic BP (DBP) was a secondary end point. The primary comparison was between participants randomly assigned to active (IND and STD) versus sham (CNTL) acupuncture. To evaluate the efficacy of acupuncture for treating hypertension, BP observations made after the initiation of antihypertensives were censored.

**TABLE 1. Treatment Definitions**

Treatment Group	Treatment Definition
IND	Corporeal acupuncture at 10 to 12 points (median: 12, counting bilateral points twice) selected and stimulated as individually prescribed. Auricular acupuncture at the 2 most active points identified with a point detector at the start of each treatment session.
STD	Corporeal acupuncture at 5 bilateral points with neutral stimulation: GB 20, LI 11, LR 03, SP 06, and ST 36. Auricular acupuncture at Heart and Jiang Ya Gou. Points selected a priori by an expert panel.
CNTL	Corporeal acupuncture at 5 bilateral points that do not fall along any TCM meridian with no stimulation. Auricular acupuncture in Darwin's tubercle and the posterior ear lobe. All corporal and auricular acupuncture points in areas considered inactive according to TCM.

The primary analysis was a linear mixed model of change in BP between randomization and weeks 2 through 10 controlling for baseline BP, use of antihypertensives in the 6 months before baseline, and gender. Treatment group, days since first treatment, and their 2-way interaction were included as fixed and random terms modeling within-participant residuals as compound symmetric over visits. Unstructured and autoregressive covariance models fit less well. Comparisons among treatment groups were tested using linear contrasts evaluating differences at week 10. Reported point estimates are least-square means.

Subgroup analyses for age, race, gender, baseline BP, history of antihypertensive use, obesity, and primary TCM diagnosis were tested by including each classification and its 2- and 3-way interactions with treatment and time in the primary model. Significance of the 3-way interaction provided evidence that treatment-specific differences in the rates of BP change varied among subgroups. Differences among treatment groups at baseline were tested by Fisher's exact test and 1-way ANOVA. Univariate changes in SBP and DBP between measurements taken immediately before and after an acupuncture treatment were tested by ANCOVA. Differences among treatment groups in the rate of antihypertensive initiation were tested by log rank test. Differences in the number of antihypertensives required were tested by the Jonckheere-Terpstra test.<sup>43</sup> Changes in laboratory values were assessed by ANCOVA. Effects on QoL were tested as the change from baseline of the Medical Outcomes Study Short Form physical and mental component scores by univariate and multivariate ANCOVA. Adequacy of masking was

tested by Fisher's exact test comparing true assignments to participant guesses, including "don't know." Confidence scores were tested by ordinal logistic regression. Differences in adverse event rates were analyzed in a negative-binomial generalized linear model. For all of the tests, significance was accepted for  $P < 0.05$  for 2-tailed tests. Analyses were performed using SAS (version 9.1.2, SAS Institute).

## Results

### Participants and Treatment

Between March 2001 and July 2002, 424 individuals provided written, informed consent and were screened for trial eligibility, of whom 192 individuals were determined eligible and interested (Figure 1). Baseline characteristics of the 192 randomly assigned participants did not differ significantly among treatment groups, except for racial profile, with more minority subjects represented in the CNTL group by chance (Table 2). Four participants lacked any follow-up data and were excluded from the primary analysis. Sample sizes are somewhat lower for data from self-administered instruments.

Of the 188 participants with follow-up, nearly all (96%) received 12 treatments over an average of 42 days (range: 35 to 65 days). IND participants received a wide range of




Figure 1. Patient flow diagram.

**TABLE 2. Participant Characteristics at Baseline**

Baseline Characteristics	IND (n=64)	STD (n=64)	CNTL (n=64)	<i>P</i>
<b>Demographics</b>				
Age, y, mean ( $\pm$ SD)	56.8 ( $\pm$ 8.4)	55.9 ( $\pm$ 10.6)	53.2 ( $\pm$ 9.5)	0.09
Male	53%	55%	55%	1.00
Hispanic or Latino	6%	0%	5%	0.16
Race				0.03
White	88%	92%	73%	
Black	11%	6%	17%	
Asian	2%	2%	9%	
<b>Clinical status and history</b>				
SBP, mean ( $\pm$ SD)	149.6 ( $\pm$ 5.8)	149.6 ( $\pm$ 7.5)	148.4 ( $\pm$ 6.3)	0.48
DBP, mean ( $\pm$ SD)	93.1 ( $\pm$ 2.9)	93.2 ( $\pm$ 2.9)	93.9 ( $\pm$ 3.4)	0.35
On antihypertensive drugs within 6 months of baseline	69%	69%	73%	0.82
Family history of hypertension	87%	83%	85%	0.84
Body mass index, mean( $\pm$ SD)	28.0 ( $\pm$ 4.8)	29.2 ( $\pm$ 4.8)	28.4 ( $\pm$ 5.2)	0.33
Smoked regularly within past 12 months	7%	8%	5%	0.93
<b>TCM primary diagnosis</b>				
Flare up of liver fire	13%	16%	17%	0.42
Liver Yang rising with kidney Yin deficiency	63%	47%	48%	
Obstruction of phlegm and dampness	19%	30%	30%	
Yin and Yang deficiency	0%	2%	3%	
Qi and blood deficiency leading to liver Yang rising	5%	6%	2%	

treatments. The most common combination of points prescribed included 3 of the 5 bilateral points used in the STD treatment (GB20, LI11, and SP06). The number of STD corporeal points prescribed for IND participants was distributed as follows: 1% received acupuncture at 0 STD points, 9% at 2 STD points, 28% at 4 STD points, 41% at 6 STD points, 20% at 8 STD points, and 2% at all 10 STD points (the sum is >100% because of rounding error). Participants received the wrong treatment on 3 occasions of 2219 treatments administered (1 CNTL participant received an IND treatment, 1 CNTL participant received a STD treatment, and 1 STD participant received an IND treatment).

### Primary and Secondary End Points

Mean SBP and DBP declined between randomization and week 10 (Figure 2). At 10 weeks, BP declines did not differ between participants randomly assigned to active (IND and STD) versus sham (CNTL) acupuncture (SBP [mean $\pm$ SE]:  $-3.56\pm 1.92$  mm Hg versus  $-3.84\pm 1.93$  mm Hg,  $P=0.90$ ; DBP:  $-4.32\pm 1.01$  mm Hg versus  $-2.81\pm 0.99$  mm Hg;  $P=0.16$ ). The 95% CIs on the estimated benefit of active versus sham acupuncture were within 4 mm Hg (active versus sham: SBP:  $-4.00$  to  $4.56$  mm Hg; DBP:  $-3.65$  to  $0.62$  mm Hg). BP declines did not differ between participants receiving IND versus STD acupuncture or IND versus CNTL acupuncture (Table 3). No treatment effects were evident after 12 months. Categorizing participants by age, race, gender, baseline BP, history of antihypertensive use, obesity, or primary TCM diagnosis did not identify any subgroups for whom the benefits of TCM acupuncture differed significantly from sham acupuncture.

On average, BP increased modestly between measurements taken before and after an acupuncture session (by 2.1/0.8 mm Hg for a participant with preacupuncture BP of 143/88 mm Hg), although individual responses were quite variable (SD:  $\pm 8.9/5.5$  mm Hg). The median interval between preacupuncture and postacupuncture measurements was 75 minutes. No significant difference was found in the immediate effect of active versus sham acupuncture (SBP [mean $\pm$ SE]:  $-0.10\pm 1.03$  mm Hg; 95% CI:  $-2.13$  to  $1.92$ ;  $P=0.92$ ; DBP:  $-1.04\pm 0.61$  mm Hg; 95% CI  $-2.24$  to  $0.15$ ;  $P=0.09$ ).

The risk of developing BP levels that required initiation of antihypertensive drugs did not differ significantly among participants randomly assigned to the 3 treatment groups ( $P=0.97$ ; Figure 3). By 10 weeks, 11% of IND participants, 19% of STD participants, and 12% of CNTL participants were on antihypertensives. The number of distinct classes of antihypertensives required to control hypertension at 10 weeks among participants using antihypertensives did not differ among the 3 treatment groups ( $P=0.51$ ). Ten-week change in physiological parameters other than BP differed significantly among treatment groups only in the case of fasting blood glucose, with a significant decrease among participants receiving sham acupuncture (IND and STD mean $\pm$ SE, CNTL mean $\pm$ SE: fasting glucose (mg/dL):  $2.33\pm 0.85$ ,  $-1.33\pm 1.23$ ;  $P=0.015$ ). The treatment groups did not differ with respect to changes in weight, blood urea nitrogen, creatinine, fasting triglycerides, total cholesterol, high-density lipoprotein, or low-density lipoprotein. STD participants reported a significant decline in mental QoL




Figure 2. SBP and DBP trajectories by treatment group.

between baseline and 10 weeks relative to IND and CNTL participants (mean  $\pm$  SE: IND:  $0.59 \pm 1.01$ ; STD:  $-2.49 \pm 0.98$ ; CNTL:  $1.17 \pm 1.09$ ;  $P=0.025$ ). Mean 10-week change in physical QoL did not differ among participants (IND:  $0.27 \pm 0.73$ ; STD:  $0.81 \pm 0.70$ ; CNTL:  $0.27 \pm 0.78$ ;  $P=0.83$ ). Overall mean change in QoL did not differ between participants receiving active versus sham acupuncture ( $P=0.35$ ).

Among the 141 participants who completed the masking assessment instrument at 4 months, the association between true random assignment and participants' guessed random assignment was nearly significant ( $P=0.06$ ). Although participants in all of the groups were equally likely to make a guess about their treatment assignment (67%;  $P=0.53$ ) and were equally confident in their guesses ( $P=0.92$ ), IND

participants were more likely to guess IND assignment (59%) than STD (19%) or CNTL (41%) participants among those guessing. Similarly, STD participants were more likely to guess STD assignment (52%) than IND (22%) or CNTL (38%) participants.

Three study-related serious adverse events occurred over 165 patient years of follow-up: 2 STD participants experienced hypertensive urgencies, and 1 CNTL participant experienced congestive heart failure. No deaths occurred. The risk of any ( $n=98$ ) or study-related ( $n=50$ ) adverse events did not differ among treatment groups ( $P>0.68$ ).


## Discussion

The results of the SHARP pilot trial did not indicate any differential benefit of active acupuncture relative to invasive

TABLE 3. BP Change From Baseline to 10 Weeks

Treatment Groups and Comparisons	N	SBP, mm Hg	DBP, mm Hg
IND	60	-3.57 (-7.33 to 0.18)	-4.67 (-6.63 to -2.70)
STD	64	-3.55 (-7.33 to 0.24)	-3.97 (-5.95 to -1.99)
CNTL	64	-3.84 (-7.62 to -0.05)	-2.81 (-4.75 to -0.87)
IND and STD vs CNTL		0.28 (-4.00 to 4.56) $P=0.90$	-1.51 (-3.65 to 0.62) $P=0.16$
IND vs STD		-0.03 (-4.90 to 4.85) $P=0.99$	-0.70 (-3.12 to 1.72) $P=0.57$
IND vs CNTL		0.27 (-4.64 to 5.18) $P=0.91$	-1.86 (-4.30 to 0.58) $P=0.14$

All values are least-square means and 95% CIs from the primary analysis.



**Figure 3.** Kaplan–Meier plot for initiation of antihypertensives after randomization. Note that criteria for initiating antihypertensives were relaxed starting at the 10-week visit. Vertical bars indicate censoring times.

sham acupuncture for controlling SBP, avoiding need for antihypertensive medications, or improving QoL or other comorbidities of hypertension, including dyslipidemia, renal function, or glucose control. No subgroups of subjects were identified that experienced a differential benefit of one treatment over another in controlling BP. The average magnitude of BP declines (SBP:  $-3.5$  to  $-3.8$  mm Hg; DBP:  $-2.8$  to  $-4.7$  mm Hg) achieved by SHARP participants was no greater than the magnitude of decrease observed in the placebo arms of 7 pharmaceutical trials analyzed by the Individual Data Analysis of Antihypertensive Intervention trials (INDANA) research committee (SBP:  $-6.8$  to  $-14.6$  mm Hg; DBP:  $-2.8$  to  $-7.2$  mm Hg; S. Pocock, unpublished data, 2003). These reductions are consistent with regression to the mean. They may also reflect changes in diet or lifestyle or other nonspecific effects of participating in clinical research. Alternatively, the observed BP reductions may represent an antihypertensive effect of invasive acupuncture unrelated to needle placement. Middlekauff et al<sup>44</sup> report that invasive sham acupuncture at a non-TCM point in the anterior deltoid resulted in a significant attenuation of the acute increase in BP induced by a mental stress test. Conversely, other researchers<sup>45,46</sup> have not observed an improvement in hemodynamics from invasive sham acupuncture.

Our results suggest that 6 weeks of twice-weekly sessions of fully individualized TCM acupuncture are unlikely to achieve clinically meaningful reductions in SBP or DBP for the average patient with mild-to-moderate hypertension relative to invasive sham acupuncture. Although the estimated 95% CIs cannot be interpreted directly as bounds on the true efficacy of active acupuncture, it is unlikely that active acupuncture reduces SBP or DBP  $>5$  mm Hg. Twice-weekly acupuncture is unlikely to be cost-effective relative to available pharmacological treatments for hypertension nor is it likely to be widely embraced by patients solely on the basis of any mild antihypertensive effect given the significant time required for acupuncture therapy. Nevertheless, it is possible that active acupuncture may yield larger benefits if the treatment is extended beyond 6 weeks. In addition, given that the diagnosing acupuncturists were prescribing holistic treat-

ments, it is possible that moderate antihypertensive effects were accompanied by other beneficial psychological or physiological effects not measured in the trial, but no benefit was seen in measures of QoL, dyslipidemia, renal function, glucose control, or weight gain.

One other well-designed randomized, controlled trial has reported a significant long-term antihypertensive effect of acupuncture.<sup>35</sup> Yin et al<sup>35</sup> report BP declines of  $-14.8/-6.9$  mm Hg in their active acupuncture group ( $n=15$ ) versus  $-4.0/-1.1$  mm Hg in the sham group ( $n=15$ ;  $P<0.05$ ) after 8 weeks of twice-weekly treatments. The study was double blind and used a noninvasive sham treatment for which masking has been validated.<sup>47</sup> At least 3 important differences exist between the SHARP pilot trial and Yin et al.<sup>35</sup> Firstly, Yin et al<sup>35</sup> used acupuncture as an adjunct to pharmaceutical management and exercise. All of the subjects studied by Yin et al<sup>35</sup> were on antihypertensives, which they continued using during the trial. Breathing and easy-walking exercises were encouraged and monitored by diary. Secondly, their participants may have been healthier. The patient sample studied by Yin et al<sup>35</sup> had lower mean baseline BP (135/83 mm Hg) and lower mean body mass index<sup>25</sup> relative to SHARP participants. Lastly and perhaps most importantly, Yin et al<sup>35</sup> compared active acupuncture to a noninvasive sham control.

What constitutes a valid acupuncture treatment and an appropriate control is a question of much debate in the field.<sup>20</sup> Indirect effects of acupuncture include remote analgesic effects of puncturing the skin with acupuncture needles<sup>48,49</sup> and psychosocial benefits of being treated by a clinician who creates an expectation of medical improvement.<sup>50</sup> Although such responses are generally regarded as placebo effects, the “intentional environment” created by acupuncturists has been identified as a component of acupuncture therapy.<sup>51</sup> In the SHARP trial, we sought to isolate the specific and direct effects of TCM acupuncture at points defined within its theory. We cannot evaluate whether needle insertion at points not along TCM-defined meridians or treatment in environments intended to foster feelings of expectation among subjects are beneficial for treating hypertension.

The SHARP trial design overcomes a number of challenges to effective evaluation of acupuncture while meeting standards of research required for evidence-based medicine. The use of separate diagnosing and treating acupuncturists and an independent vascular medicine team permitted effective masking of treatment randomizations and BP responses. Inclusion of both IND and STD treatment groups permitted evaluation of any differential benefit of acupuncture prescriptions tailored to a subject’s specific presentation. The use of clear eligibility criteria, objective randomization, careful BP monitoring, specific guidelines for use of antihypertensives, and rigorous analysis raise the standards for research into acupuncture and other therapies used in complementary and alternative medicine.

A number of factors limited interpretations from the SHARP trial. Although larger than any previous study, the SHARP pilot trial was not designed to detect small effects. The conclusion of no active acupuncture effect would be stronger if the CIs of the estimated differences between treatments were narrower. Use of

acupuncture without complementary use of antihypertensives does not reflect the usual practice among acupuncturists for subjects with stage 2 hypertension. Twelve treatment sessions spread over 6 to 8 weeks may be inadequate to reduce systolic hypertension reliably. The small magnitude of any immediate antihypertensive effect of active acupuncture may suggest that acupuncture treatments were not adequately administered given that most published work<sup>30,38,45,46</sup> and practicing acupuncturists report sizeable reductions in both SBP and DBP immediately after treatment. Lastly, use of an invasive sham control group prevented estimation of nonspecific effects of acupuncture, which may constitute an important component of clinical acupuncture.

### Perspectives

Twelve sessions of TCM acupuncture do not appear to control hypertension better than invasive sham acupuncture when used as monotherapy. Some participants experienced substantial improvements in their hypertension, but no subgroups of participants could be identified from baseline characteristics that experienced a differential benefit of active acupuncture. Further research is required to determine whether acupuncture can enhance clinical management of hypertension if used in combination with antihypertensive agents, over longer periods, or among specific subgroups of patients.

### Acknowledgments

Thanks are expressed to the participants of the SHARP trial for their time and commitment, to our acupuncture consultants (Jian Kong, Jing Liu, Weidong Lu, and Zhen-zhen Zhang); the staff at the Hypertension Center (Sharon Maginnis, Jean Sullivan, and Mildred Wasileski), the Acupuncture Clinic (Kate Billings, Edward Chiu, Christian Connors, Amy Hull, Diane Iuliano, Zhi Ping Li, and Nicole Stockholm), and the Data Coordinating Center (Patricia Connell, Allison Gemmel, and Julie Miller); to Chang Shik Yin and Yun-ju Kim for sharing prepublished manuscripts and bibliographic suggestions; and to Stuart Pocock for sharing data from the INDANA database.

### Source of Funding

This work was supported by Cooperative Agreement U01 AT00210, awarded to New England Research Institutes, Inc, by the National Center for Complementary and Alternative Medicine, National Institutes of Health.

### Disclosures

None.

### References

- Mensah GA. The global burden of hypertension: good news and bad news. *Cardiol Clin*. 2002;20:181–185.
- Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJ. Selected major risk factors and global and regional burden of disease. *Lancet*. 2002;360:1347–1360.
- Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jr., Jones DW, Materson BJ, Oparil S, Wright JT, Jr, Roccella EJ. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003;42:1206–1252.
- Tindle HA, Davis RB, Phillips RS, Eisenberg DM. Trends in use of complementary and alternative medicine by US adults: 1997–2002. *Altern Ther Health Med*. 2005;11:42–49.
- Lin MC, Nahin R, Gershwin ME, Longhurst JC, Wu KK. State of complementary and alternative medicine in cardiovascular, lung, and blood research: executive summary of a workshop. *Circulation*. 2001;103:2038–2041.
- Kaptchuk TJ. *The Web That Has No Weaver: Understanding Chinese Medicine*. Revised ed. Chicago, IL: Contemporary Books; 2000.
- Xu X. Acupuncture in an outpatient clinic in China: a comparison with the use of acupuncture in North America. *South Med J*. 2001;94:813–816.
- Napadow V, Kaptchuk TJ. Patient characteristics for outpatient acupuncture in Beijing, China. *J Altern Complement Med*. 2004;10:565–572.
- Flaws B, Sionneau P. *The Treatment of Modern Western Medical Diseases With Chinese Medicine: A Textbook & Clinical Manual*. 1st ed. Boulder, CO: Blue Poppy Press; 2001.
- Ernst E, Pittler MH. The effectiveness of acupuncture in treating acute dental pain: a systematic review. *Br Dent J*. 1998;184:443–447.
- Vickers AJ. Can acupuncture have specific effects on health? A systematic review of acupuncture antiemesis trials. *J R Soc Med*. 1996;89:303–311.
- Lee A, Done ML. The use of nonpharmacologic techniques to prevent postoperative nausea and vomiting: a meta-analysis. *Anesth Analg*. 1999;88:1362–1369.
- White A. A cumulative review of the range and incidence of significant adverse events associated with acupuncture. *Acupunct Med*. 2004;22:122–133.
- Smith CA, Collins CT, Cyna AM, Crowther CA. Complementary and alternative therapies for pain management in labour. *Cochrane Database Syst Rev*. 2003;2:CD003521.
- Soeken KL. Selected CAM therapies for arthritis-related pain: the evidence from systematic reviews. *Clin J Pain*. 2004;20:13–18.
- Trinh KV, Phillips SD, Ho E, Damsma K. Acupuncture for the alleviation of lateral epicondyle pain: a systematic review. *Rheumatology (Oxford)*. 2004;43:1085–1090.
- Vickers AJ, Rees RW, Zollman CE, McCarney R, Smith CM, Ellis N, Fisher P, Van Haselen R. Acupuncture for chronic headache in primary care: large, pragmatic, randomised trial. *BMJ*. 2004;328:744.
- Berman BM, Lao L, Langenberg P, Lee WL, Gilpin AM, Hochberg MC. Effectiveness of acupuncture as adjunctive therapy in osteoarthritis of the knee: a randomized, controlled trial. *Ann Intern Med*. 2004;141:901–910.
- Linde M, Fjell A, Carlsson J, Dahlof C. Role of the needling per se in acupuncture as prophylaxis for menstrually related migraine: a randomized placebo-controlled study. *Cephalalgia*. 2005;25:41–47.
- Birch S. A review and analysis of placebo treatments, placebo effects, and placebo controls in trials of medical procedures when sham is not inert. *J Altern Complement Med*. 2006;12:303–310.
- Anshelevich Iu V, Merson MA, Afanas'eva GA. Serum aldosterone level in patients with hypertension during treatment by acupuncture [in Russian]. *Ter Arkh*. 1985;57:42–45.
- Huang H, Liang S. Acupuncture at otaocupoint heart for treatment of vascular hypertension. *J Tradit Chin Med*. 1992;12:133–136.
- Akhmedov TI, Vasil'ev Iu M, Masliaeva LV. The hemodynamic and neurohumoral correlates of the changes in the status of hypertension patients under the influence of acupuncture [in Russian]. *Ter Arkh*. 1993;65:22–24.
- Zhou Y, Wang Y, Fang Z, Xia C, Liu B, Chen Q, Zhang F. Influence of acupuncture on blood pressure, contents of NE, DA and 5-HT of SHR and the interrelation between blood pressure and whole blood viscosity [in Chinese]. *Zhen Ci Yan Jiu*. 1995;20:55–61.
- Chiu YJ, Chi A, Reid IA. Cardiovascular and endocrine effects of acupuncture in hypertensive patients. *Clin Exp Hypertens*. 1997;19:1047–1063.
- Chao DM, Shen LL, Tjen ALS, Pitsillides KF, Li P, Longhurst JC. Naloxone reverses inhibitory effect of electroacupuncture on sympathetic cardiovascular reflex responses. *Am J Physiol*. 1999;276:H2127–H2134.
- Li P, Tjen ALS, Longhurst JC. Rostral ventrolateral medullary opioid receptor subtypes in the inhibitory effect of electroacupuncture on reflex autonomic response in cats. *Auton Neurosci*. 2001;89:38–47.
- Tjen-A-Looi SC, Li P, Longhurst JC. Prolonged inhibition of rostral ventral lateral medullary premotor sympathetic neurons by electroacupuncture in cats. *Auton Neurosci*. 2003;106:119–131.
- Zhou W, Fu LW, Tjen ALSC, Li P, Longhurst JC. Afferent mechanisms underlying stimulation modality-related modulation of acupuncture-related cardiovascular responses. *J Appl Physiol*. 2005;98:872–880.
- Tam KC, Yiu HH. The effect of acupuncture on essential hypertension. *Am J Chin Med*. 1975;3:369–375.
- Dong J. Acupuncture treatment of hypertension: a report of 70 cases. *Int J Clin Acupunct*. 1996;7:173–175.

32. Guo W, Ni G. The effects of acupuncture on blood pressure in different patients. *J Tradit Chin Med.* 2003;23:49–50.
33. Yu Y, Tang L. Investigation into the potential synergism between body and auricular acupuncture points for the treatment of hypertension [letter]. *Am J Acupunct.* 1995;23:184–185.
34. Lu B, Ren S, Hu X, Lichstein E. A randomized controlled trial of acupuncture and acupressure treatment for essential hypertension [abstract]. *Am J Hypertens.* 2000;13:S185.
35. Yin CS, Seo BK, Park HJ, Lim S, Cho M, Jung WS, Choue R, Kim CH, Lee H, Koh HG. Acupuncture, a promising adjunctive therapy for essential hypertension: a double-blind, randomized, controlled trial. *Neurological Research.* In press.
36. Kraft K, Coulon S. Effect of a standardized acupuncture treatment on complaints, blood pressure and serum lipids of hypertensive, postmenopausal women. A randomized, controlled clinical study [in German]. *Forsch Komplementarmed.* 1999;6:74–79.
37. Jiang X. Effects of magnetic needle acupuncture on blood pressure and plasma ET-1 level in the patient of hypertension. *J Tradit Chin Med.* 2003;23:290–291.
38. Robinson RC, Wang Z, Victor RG, Vongpatanasin W, Zhang W, Abbas A, Gatchel RJ, Kaplan NM. Lack of effect of repetitive acupuncture on clinic and ambulatory blood pressure [abstract]. *Am J Hypertens.* 2004;17:S33.
39. Kalish LA, Buczynski B, Connell P, Gemmel A, Goertz C, Macklin EA, Pian-Smith M, Stevens S, Thompson J, Valaskatgis P, Wayne PM, Zusman RM. Stop Hypertension with the Acupuncture Research Program (SHARP): clinical trial design and screening results. *Control Clin Trials.* 2004;25:76–103.
40. Cheng X, ed. *Chinese Acupuncture and Moxibustion.* Revised ed. Beijing, China: Foreign Language Press; 1999.
41. Ware JE, Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care.* 1992;30:473–483.
42. International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use. Clinical safety data management: Definitions and standards for reporting. *Federal Register* 1995;60:11284–11287.
43. Hollander M, Wolfe DA. *Nonparametric Statistical Methods.* New York, NY: John Wiley & Sons; 1973.
44. Middlekauff HR, Yu JL, Hui K. Acupuncture effects on reflex responses to mental stress in humans. *Am J Physiol Regul Integr Comp Physiol.* 2001;280:R1462–R1468.
45. Williams T, Mueller K, Cornwall MW. Effect of acupuncture-point stimulation on diastolic blood pressure in hypertensive subjects: a preliminary study. *Phys Ther.* 1991;71:523–529.
46. Li P, Ayannusi O, Reid C, Longhurst JC. Inhibitory effect of electroacupuncture (EA) on the pressor response induced by exercise stress. *Clin Auton Res.* 2004;14:182–188.
47. Park J, White A, Stevinson C, Ernst E, James M. Validating a new non-penetrating sham acupuncture device: two randomised controlled trials. *Acupunct Med.* 2002;20:168–174.
48. Le Bars D, Dickenson AH, Besson JM. Diffuse noxious inhibitory controls (DNIC). II. Lack of effect on non-convergent neurones, supraspinal involvement and theoretical implications. *Pain.* 1979;6:305–327.
49. Xu WD, Zhu B, Rong PJ, Bei H, Gao XY, Li YQ. The pain-relieving effects induced by electroacupuncture with different intensities at homotopic and heterotopic acupoints in humans. *Am J Chin Med.* 2003;31:791–802.
50. Kirsch I. *How Expectancies Shape Experience.* 1st ed. Washington, DC: American Psychological Association; 1999.
51. Kaptchuk TJ. The placebo effect in alternative medicine: can the performance of a healing ritual have clinical significance? *Ann Intern Med.* 2002;136:817–825.